



# Successful Treatment of Borreliosis



Borrelia infections following tick bites are by no means uncommon. The pathogens *Borrelia burgdorferi* or *Borrelia burgdorferi* (Lyme borreliosis) are gram-negative group of bacteria and among other things, can cause recurrent fever with sickness as well as pins in the head, limbs, joints and back. The disease can be diagnosed reliably by a blood test. In a biological practice, the following case was treated successfully which primarily may be ascribed to the effect of SANUM preparations.

In September 1997 an elevated IgM count following a tick bite was determined in the 49-year-old female patient. She was suffering from strong pains in the joints - particularly the hip joint - and was

very weak, and she also had an itching rash on her face, elbows and stomach, in the liver region and on her shins. First of all an animal protein-free diet was recommended. Treatment started with ALKALAN and Gastrohelp, and after three weeks UTILIN „S“ 6X capsules (one capsule every two weeks) and SANUKEHL BRUCEL 6X (1 ampoule twice weekly) were administered additionally.

The patient began to improve steadily during this course of treatment, and the joint pains and skin became less and less. Six weeks later, the borrelia titer was checked and proved negative. As the doctor, who had previously been consulted by the patient doubted these results, her blood was tested once

more in a different laboratory, but they found no trace of the borrelia titer either. Two months later, the patient came to the practice once more: she was still improving, she had regained her original strength, and the joint pains and the very troublesome skin rash on various parts of her body had disappeared and not returned.

**First published in the German language in the SANUM-Post magazine (47/1999)**

**© Copyright 1999 by Semmelweis-Institut GmbH, 27318 Hoya (Weser), Germany**

**All Rights Reserved**