



The Therapy of Thrombocytosis with SANUM Remedies

by

Dr. med. Thomas Rau



That thrombocytes have a connection with the mucor racemosus series is difficult to understand for allopathically trained people, just as it is difficult to understand that the protein fibrinogen contains endobionts of a much lower valence. Yet the therapeutic results confirm this way of thinking. It is therefore necessary to administer MUCOKEHL to treat thrombocytæmia or thrombocytosis. MUCOKEHL suppositories prove to be most effective, as their effect on medulla or spleen is better. It is just as important to ensure that the valences of a lower developmental state are kept at that low level. This is best done with MUCOKEHL Excretion 6X, 1 ampoule twice a week or every other day 10 drops taken nasally. Furthermore the administration of SANUVIS is also required, here 3 x 20 drops a day would be the minimum. The further decrease of the developmental state is ensured by the application of base preparations, such as ALKALAN,

2 x 1/2 tsp a day and mainly by a richly alkaline diet.

Thrombocytoses are also always connected with an excessive spleen activity and have thus to be treated with high doses of vitamin C, 2–3 grams a day, folic acid and vitamin B-complex. You have to remember that this lowers the high valences, turning them towards fibrinogen or hypercoagulability. This has to be counteracted with colonic irrigation, a strictly vegetarian diet and the exclusion of all dairy products. Enzymes, i. e. Wobenzyme, 3 x 4 coated tablets taken between meals with a lot of liquid, are a successful supplement and also quite important. Kreon (Solvay), a pancreas medicament (2 x 1 capsules) can be supplemented as well.

The therapy of thrombocytosis is always a long-term commitment and must be supported homoeopathically. Lachesis is very frequently used, Luesinum less so. Of course,

any therapy is individually specified according to the dark-field results.

The macrophage activity supports the therapy with its protein dissolving effect and can be increased immune-biologically with the administration of UTILIN 6X and RECARCIN 6X capsules, taken once a week.

Dr. med Thomas Rau

First published in the German language in the Sanum-Post magazine (62/2003)

**© Copyright by Semmelweis-Institut GmbH, 27318 Hoya, Germany
All Rights Reserved.**