



Illnesses of the Ear

by Harald Krebs



EAR ECZEMAS

causes:

Mechanical or chemical irritation
Allergies
Metabolic maladies
Excretory disorders of the liver, intestines or kidneys

Symptoms:

Typical signs of inflammation
Redness
Swelling
Fever
Pain
Vesicle pustule and rhagade formation

Advisable:

Perform a lab test, such as a hemogram, especially IgE level, if indicated; also investigate liver parameters.

Therapy recommendation:

Injection therapy:

Every 14 days, 1 Ampule Latensin 6X i.m.
Intercalation of injections for intensive diversion:

- Sanukehl Cand 5X i.m.
- Sanukehl Myc 5X i.m.
- Sanukehl Strep 5X i.m.

The Sanukehl preparations (Haptenes) must always be injected by themselves, i.e. they are administered in the week in which no Latensin injection is administered. Begin with the injection of Sanukehl Cand 5X, after 14 days Sanukehl Myc 5X, and after 14 more days Sanukehl Strep 5X.

Important:

When Sanum Haptenes are administered, the situation can temporarily worsen at first. It is therefore absolutely necessary, during injection treatment, to make sure that the patient drink sufficient quantities of fluids.

Oral medication:

Eczema at the orifice of the ear: Silicea D12 tablets:
1 tablet twice daily

Eczema in the auditory canal: Petroleum 12X dil.: 10 drops twice daily

Eczema with vesicle formation: Sepia 6X tablets: 1 tablet 3 times daily

The following tea mixture has proven effective for diversion:

Take: (as per J. Karl)

Fumitory herb 10g
Stip. Dulcamarae 10g
Couch gras radix 10g
Linden flowers 40g
Stinging nettle herb 30g
Mix according to specification: 2 tsp. with 2 cups of boiling water, the tea must stand for 10 minutes. Drink 3 cups of this tea, daily for a period of 3-4 weeks

EAR MYCOSIS

Causes:

Mainly from swimming in swimming pools

Symptoms:

Itching, scaly, whitish-yellowish deposits in the outer ear passage

Advisable:

Confirm the diagnosis by taking a smear and culturing it.

Therapy recommendation:

Injection therapy:

Twice weekly compound injection i.m.
1 Ampule Pefrakehl 6X
1 Ampule Sanuvis

Oral medication:

Pefrakehl 5X drops: once daily 10 drops before breakfast
Phoenix detoxification therapy:
Phoenix Phoenhepan: 3 times daily 60 drops for 3 days.
Phoenix Solidago: 3 times daily 60 drops for 3 days.
Phoenix Antitox: 3 times daily 10 drops for 3 days.
Now repeat the Phoenix detoxification cycle for a period of 45 days.

In very therapy-resistant cases, the administration of suppositories in the evening is advised:

Mon Nigersan 3X suppository
Tue Pefrakehl 3X suppository
Wed Pefrakehl 3X suppository
Thu Pefrakehl 3X suppository
Fri Nigersan 3X suppository

Depending on the results with the above therapy, this regimen can be repeated for one more week.



ACUTE HEARING LOSS

Causes:

One is dealing here with an inner-ear lesion of indeterminate etiology. One thing is certain: a microcirculation disturbance has triggered the hearing loss. Autoimmune processes in the body may also be a factor.

This is an emergency situation.

Symptoms:

Sensation of pressure in the ear

Increasing tinnitus

Sudden onset of considerable hearing loss, including total deafness in extreme cases

Advisable:

Consult a specialist or clinic immediately

Therapy recommendation:

Injection therapy:

First Aid

Carbo capsule (Heel) immediately 1 Ampule i.v.

Compound injection strictly i.c. at the mastoid process:

1 Ampule Mucokehl 5X

1 Ampule Lidocaine

Oral medication:

Mucokehl 4Xcapsules

2 capsules before breakfast and 1 capsule before going to bed

Important:

Should noises in the ear continue after specialist's treatment, try out the following therapy:

2 x weekly compound injection strictly i.c. at the mastoid process:

1 Ampule Mucokehl 5X

1 Ampule Lidocaine

At the Same time administer an infusion of 100 ml NaCl solution

5 Ampules Ginkgobakehl 4X

First published in the English language in the EXPLORE magazine (Vol.9#2 1999)

©Copyright by

Semmelweis Institute GmbH

27318 Hoya, (Weser), Germany

All Rights Reserved